



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Multi-network-based diffusion analysis reveals vertical cultural transmission of sponge tool use within dolphin matriline

Wild, Sonja ; Allen, Simon J ; Krützen, Michael ; King, Stephanie L ; Gerber, Livia ; Hoppitt, William J E

Abstract: Behavioural differences among social groups can arise from differing ecological conditions, genetic predispositions and/or social learning. In the past, social learning has typically been inferred as responsible for the spread of behaviour by the exclusion of ecological and genetic factors. This 'method of exclusion' was used to infer that 'sponging', a foraging behaviour involving tool use in the bottlenose dolphin (*Tursiops aduncus*) population in Shark Bay, Western Australia, was socially transmitted. However, previous studies were limited in that they never fully accounted for alternative factors, and that social learning, ecology and genetics are not mutually exclusive in causing behavioural variation. Here, we quantified the importance of social learning on the diffusion of sponging, for the first time explicitly accounting for ecological and genetic factors, using a multi-network version of 'network-based diffusion analysis'. Our results provide compelling support for previous findings that sponging is vertically socially transmitted from mother to (primarily female) offspring. This research illustrates the utility of social network analysis in elucidating the explanatory mechanisms behind the transmission of behaviour in wild animal populations.

DOI: <https://doi.org/10.1098/rsbl.2019.0227>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-184389>

Journal Article

Accepted Version

Originally published at:

Wild, Sonja; Allen, Simon J; Krützen, Michael; King, Stephanie L; Gerber, Livia; Hoppitt, William J E (2019). Multi-network-based diffusion analysis reveals vertical cultural transmission of sponge tool use within dolphin matriline. *Biology Letters*, 15(7):20190227.

DOI: <https://doi.org/10.1098/rsbl.2019.0227>

Multi-Network-Based Diffusion Analysis reveals vertical cultural transmission of sponge tool use within dolphin matriline

Sonja Wild^{1,2*}; Simon J. Allen^{2,3,4}; Michael Krützen²; Stephanie L. King^{3,4}; Livia Gerber²; William J.E. Hoppitt^{1,5}

1. School of Biology, University of Leeds, Leeds, United Kingdom

2. Evolutionary Genetics Group, Department of Anthropology, University of Zurich, Zurich, Switzerland

3. School of Biological Sciences, University of Bristol, Bristol, United Kingdom

4. School of Biological Sciences, Oceans Institute, University of Western Australia, Perth, Australia

5. School of Biological Sciences, Royal Holloway, University of London, United Kingdom

*Corresponding author: sonja.wild@uzh.ch

Keywords:

bottlenose dolphins; vertical social learning; ecology; genetics; culture, network-based diffusion analysis, NBDA; tool use; sponging

Abstract

Behavioural differences among social groups can arise from differing ecological conditions, genetic predispositions and/or social learning. In the past, social learning has typically been inferred as responsible for the spread of behaviour by the exclusion of ecological and genetic factors. This 'method of exclusion' was used to infer that 'sponging', a foraging behaviour involving tool use in the bottlenose dolphin (*Tursiops aduncus*) population in Shark Bay, Western Australia, was socially transmitted. However, previous studies were limited in that they never fully accounted for alternative factors, and that social learning, ecology and genetics are not mutually exclusive in causing behavioural variation. Here, we quantified the importance of social learning on the diffusion of sponging, for the first time explicitly accounting for ecological and genetic factors, using a multi-network version of 'network-based diffusion analysis' (NBDA). Our results provide compelling support for previous findings that sponging is vertically socially transmitted from mother to (primarily female) offspring. This research illustrates the utility of social network analysis in elucidating the explanatory mechanisms behind the transmission of behaviour in wild animal populations.

Introduction

Various mechanisms can be responsible for causing behavioural differences among social groups or populations [1]. The *cultural hypothesis* states that behavioural variation is the result of social transmission of different behavioural innovations. The *ecological hypothesis*, on the other hand, proposes that behavioural differences among groups can be attributed to differing ecological conditions. Finally, the *genetic hypothesis* assumes that different groups are genetically predisposed to behave in different ways [1].

The last few decades have seen increasing interest in animal cultural phenomena, *i.e.*, behaviours that are socially transmitted among conspecifics [1]. Various methods have been used to identify social learning in animal populations. For example, the *method of exclusion* (also termed *group contrast method*, or *ethnographic method*) – commonly used among primatologists in the past *e.g.* [2,3] - identifies patterns of variation in the behavioural repertoires of the population in question and infers social transmission as at least partly responsible for differing behaviours by excluding genetic and ecological differences as sufficient explanations [4] [p. 132].

The method of exclusion has also been used to assess patterns of transmission of ‘sponging’, a foraging behaviour involving tool use in a population of Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) in Shark Bay, Western Australia [5]. This behaviour involves dolphins carrying conical sponges as protective ‘gloves’ on their rostra when foraging for buried prey [6]. Sponging is female-biased, and almost all sponging dolphins possess the same mitochondrial haplotype, *i.e.*, belong to the same matriline [5,7]. As the deep-water channels where sponging occurs were used by both ‘spongers’ and ‘non-spongers’, a purely ecological explanation seemed unlikely [5]. By considering 10 different pathways of potential genetic inheritance (x-linked and autosomal), Krützen *et al.* [5] inferred that sponging was vertically socially transmitted from mother to female offspring.

The method of exclusion has been criticised, however, with considerable debate over its utility [8–10]. Laland and Janik [9] argued that it is impossible to take all plausible explanations for the spread of behaviour into account, and therefore, that social learning can never be inferred with absolute certainty, leading to increased rates of false claims of culture [4]. Furthermore, they argued that social learning, ecology and genetics are not necessarily mutually exclusive [9,10]. Instead, they can simultaneously shape behaviour in a population, warranting a more nuanced approach to disentangle the relative contributions of the three drivers of behavioural variation.

In an attempt to resolve the animal cultures debate, more quantitative methods to infer social learning have been developed. For example, using repertoire-based methods on long-term

behavioural data from eleven orang-utan (*Pongo* spp.) populations, Krützen and colleagues [11] showed that neither uniquely genetic nor ecological components explained the total observed variance with regards to putative cultural elements, corroborating a cultural explanation. Further, ‘network-based diffusion analysis’ (NBDA) [12,13], a network-based approach allowing the quantification of the importance of social learning on the spread of behaviour, has been used increasingly in recent years to detect and quantify social learning in animal populations, e.g. [14,15]. NBDA infers social transmission if the spread of a behaviour follows the social network, assuming that more closely associated individuals have more opportunities to learn from each other [13,16]. Multi-network NBDA allows the inclusion of several different networks to quantify the relative importance of transmission along different pathways [17].

Here, we use multi-network NBDA to quantify the relative importance of social learning, ecological factors and genetic relatedness on the spread of sponge tool use in the dolphin population of Shark Bay, Western Australia. Furthermore, we distinguished between different pathways of social learning, namely vertical (between mother and offspring) and horizontal/oblique learning (among peers/between older and younger generations, respectively).

Methods:

Field methods

We collected association and behavioural data during boat-based surveys using standardised sampling methods for cetaceans between 2007 and 2018 in the western gulf of Shark Bay, Western Australia. On approach to each dolphin group, we recorded GPS location, determined group composition during the first five minutes of each encounter using long-established photo-identification techniques [18], and recorded predominant group behaviour. All occurrences of sponging were recorded and an individual was deemed a ‘sponger’ once it had been seen carrying a sponge on at least two independent occasions. Biopsy samples were taken on an opportunistic basis using a system designed specifically for sampling cetaceans [19].

Genetic methods

To test for a genetic predisposition for developing sponging behaviour, we obtained a measure of genetic biparental relatedness for each dyad. Individuals for which biopsies were available (N = 295) were genetically sexed [20] and genotypes determined based on 27 microsatellite markers (SI, Tab. 1). Using COANCESTRY 1.0.1.7 [21], we calculated dyadic biparental relatedness based on genotypes for individuals with no more than three microsatellite loci missing (N = 293), using the estimator TrioML [22] (SI). With a cut-off point of seven sightings (see below), genetic data were available on

226 out of 415 individuals, resulting in 25,425 unique dyads. For the remaining 189 individuals where no genetic information was available (60,480 dyads) we used the population average relatedness of 0.043.

We also statistically controlled for a correlation between matriline membership and sponging behaviour by sequencing a 468 bp-long fragment of the mitochondrial DNA (mtDNA) control region ('d-loop') to assign dolphins to mtDNA haplotypes [23].

Network constructions and NBDA

To assess the relative importance of social learning, ecological factors and genetics in promoting the spread of sponging, we ran multi-network NBDA [17] using four different networks (NBDA package v0.6.1 [24] in R 3.5.1 [25]). The first social network assessed vertical learning between mother and offspring, with entries of 1 between mother and known offspring and all other connections set to 0. We created the network based on behaviourally and genetically identified mother-offspring pairs (N = 294; SI). The second social network allowed for horizontal/oblique (henceforth 'horizontal') learning using dyadic association strengths (Simple Ratio Index [26]) among all individuals but excluding mother-offspring associations, which were set to 0. Sightings of the same or a subset of the original group within two hours were excluded. Association matrices were created using R package 'asnipe' [27]. The third, ecological network contained dyadic home range overlap as a proxy of the environmental similarity experienced by individuals. We created home ranges using individual GPS locations based on 95% Epanechnikov kernel density estimates ('adehabitatHR' [28]) with a customized smoothing factor (SI). Dyadic home range overlap (95%) was calculated using the 'utilization distribution overlap index' ('adehabitatHR') [28,29]. Finally, the fourth network contained measures of dyadic biparental genetic relatedness among individuals. Since NBDA infers social learning if a behaviour follows the social network, there is a trade-off between sample size and data quality. Dropping individuals with few sightings can increase certainty about the strengths of connections but, at the same time, decrease power of NBDA to reliably detect social learning if linking individuals are removed [30]. We ran a simulation to select a threshold that maximises power of NBDA to detect social learning, revealing maximum power at seven sightings (SI) [30]. In all networks, we therefore only considered individuals with a minimum of seven observations.

We then applied the 'order-of acquisition diffusion analysis' (ODA) variant of NBDA [13] (SI). For several individuals, the order of acquisition of sponging was unknown, as they were likely already spongers when first encountered. In NBDA models, such individuals can be taken to be 'informed' at the start of the diffusion (termed 'demonstrators') [13]. We considered all individuals

as demonstrators who had been seen carrying a sponge within the first two encounters where predominant group behaviour was foraging. We argue that an individual's information state can be determined with reasonable certainty after two sightings, given spongers carry sponges 96% of the time when foraging [31]. Maternity data were unavailable for nine individuals who acquired sponging after 2007. These nine individuals were excluded as learners, but we allowed for other individuals having learned from these spongers (SI).

We included several individual-level variables (ILVs) with potential influence on the learning rate: sex; average water depth of each individual's sightings (a proxy for habitat use, since sponging occurs in deep-water channels [32]); average group size (since sponging is a solitary activity [31]), and mitochondrial haplotype as a reduced two-level factor (either haplotype E (=sponging haplotype in the western gulf [7]), or other) to avoid overfitting of models. Sex was determined genetically and/or by the presence of a dependent calf for females. In an NBDA, the strength of transmission through a network ('s parameter') is estimated relative to a baseline rate of asocial learning. This baseline was set to the mean of all continuous variables, at the mid-point between males and females, and haplotype E (set as the reference level for this factor).


We fitted OADA with and without transmission through the networks and with all possible combinations of networks and ILVs [13]. Thereby, ILVs were allowed to influence both social and asocial learning rates independently ('unconstrained' models [4]; SI). Support for each model was calculated based on the Akaike Information Criterion corrected for sample size (AICc) [33]. To provide a more robust inference about strength of transmission for the different networks and the influence of ILVs, model averaging methods were employed [33]. We calculated 95% confidence intervals for model parameters using the profile likelihood method, conditional on the best performing model (SI).

Results


Between 2007 and 2018, 5,300 dolphin groups were encountered in the western gulf of Shark Bay and >1,000 different dolphins identified (Fig. 1a). Sponging was observed on 825 occasions and restricted to the deep-water channels within the study area (Fig. 1b). A total of 76 individuals were identified as spongers, of which 49 were confirmed female, 14 male and 13 of unknown sex.

After removal of individuals with fewer than seven sightings, as well as eight offspring that were either dependent calves at the time of analyses or had died before weaning, 415 individuals remained, of which 62 were spongers (18 learners, of which 9 were removed due to missing maternity data, and 44 demonstrators). All spongers with maternity data available were born to

sponging mothers. All spongers with genetic data available carried haplotype E, with one exception:
a male sponger with haplotype H (but see SI).



Locations of a) all dolphin groups encountered in the western gulf of Shark Bay between 2007 and 2018; and b) observations of sponging behaviour, which primarily occurred in deep (>10 m) water channels (white areas).



Multi-network NBDA revealed most support (based on Akaike weights) for transmission of sponging through the vertical social network, while transmission through the horizontal, ecological and relatedness networks (or combinations thereof) received little support.

Multi-network NBDA revealed most support for models with transmission through the vertical social network ($\sum w_i = 0.837$), while asocial learning, and transmission through the horizontal, ecological or genetic network (or any combination of the four networks) received much less support ($\sum w_i < 0.1$; Fig. 2). In the best performing model, which included vertical social transmission and sex influencing social learning, s (the rate of social transmission from mothers to offspring) was estimated to be 1.23×10^{10} times greater than the rate of asocial learning (95% C.I. [33.1; infinity]; $\sum w_i = 0.425$). The social learning rate was an estimated 126 times higher for females than males (95% C.I. [9.5; 2897]; $\sum w_i = 0.975$). This corresponds to approximately 100% of spongers learning sponging socially from

their mothers (95% C.I. [98.9; 100]). Average group size, average water depth and haplotype did not influence social or asocial learning rate (all $\sum w_i < 0.5$; SI).

Discussion

We applied multi-network NBDA to sponging behaviour, revealing overwhelming support for social transmission through the vertical mother-offspring network, with little or no support for transmission through the horizontal association, ecological or genetic networks. Moreover, despite the restriction of sponging to channel habitat [32,34], our analysis suggests that ecological factors play only a minor role once vertical social learning has been taken into account.

Low support for transmission through the genetic network confirms previous findings that sponging individuals in the western gulf are not more closely related than expected by chance [7]. This stands in contrast to findings from the eastern gulf of Shark Bay, where spongers show higher relatedness than the population average, suggesting a more recent common ancestry [5].

We further confirm a previously documented female sex-bias [7,31,35], which is presumably due to differing sex-specific reproductive strategies between males and females [31]. After weaning, male dolphins must focus on forming multi-male alliances to coerce and consort oestrous females [36–38]. This requires significant investment in social relationships and is, therefore, largely incompatible with a time-consuming, solitary and difficult-to-master activity like sponging [31,39]. Meanwhile, female offspring are expected to invest more into developing foraging skills to maximize food intake compared to male offspring [40,41]. Alternatively, Zefferman [42] proposed that the female sex-bias could be the result of a maternal teaching strategy, arguing that teaching a daughter would result in higher long-term fitness for a female: a potential advantage of sponging for a son would last only one generation, while a daughter can pass on the behaviour to subsequent generations which all gain potential benefits associated with sponging. Just 22% of spongers with known sex in the western gulf were males, which corresponded to previously suggested proportions of male offspring learning sponging from their mothers in Shark Bay's eastern gulf [31, but see 43].

Given haplotype similarity among spongers, some researchers have argued that mitochondrial genes themselves might predispose dolphins to learning the sponging behaviour [9]. However, we find no evidence that being a member of a particular mtDNA matriline has an effect on the rate at which dolphins learn sponging, as per previous research [44]. Our findings instead support the hypothesis that maternal vertical transmission of both the sponging behaviour and mtDNA results in haplotype similarity among spongers, a phenomenon referred to as 'cultural hitchhiking' - a form of gene-

culture co-evolution in which a neutral genetic locus is inherited in parallel with a matrilineally transmitted cultural behaviour [45].

McElreath and Strimling's [46] mathematical models predict the conditions for the evolution of purely vertical transmission, concluding that "neither [vertical nor oblique] transmission should be expected to dominate the other across all domains" [46]. Sponging is just one foraging strategy exhibited by the dolphins, and other strategies may be transmitted obliquely and horizontally. Following McElreath and Strimling's models [46], we suggest that sponging is transmitted vertically either because the relevant environment (e.g. availability of sponges) may be stable. Alternatively, it may only be possible for a dolphin to learn sponging from its mother, if, for example, it requires repeated observations from close quarters.

The application of multi-network NBDA to sponging behaviour in the dolphins of western Shark Bay allowed us to quantify the effects of social learning on behaviour, whilst explicitly accounting for the influence of ecological and genetic factors for the first time. Documenting a strong effect of vertical social learning from mother to offspring, our findings provide strong quantitative evidence to support the claim that sponging is a case of vertically transmitted culture in the bottlenose dolphins of Shark Bay [5].

Acknowledgements

We thank Shark Bay Resources and the Useless Loop community for logistic support, and all researchers and volunteers who have contributed to data collection for the Dolphin Innovation Project.

Author contributions

Conceptualization - SW, MK, SJA and WH. Funding – MK, SLK. Data collection - SW, SJA, MK, LG, SLK. Formal analysis – SW, WH. Writing - all. All authors approved of, and agreed to be held accountable for, the final manuscript.

Accessibility

Code is available in ESM, and data on dryad [47]:
<https://datadryad.org/review?doi=doi:10.5061/dryad.sc26m6c>.

Funding

This research was funded by: Swiss National Science Foundation (31003A_149956), Seaworld Research & Rescue Foundation Inc. (SWRRFI), National Geographic Society, A.H. Schultz Stiftung, Claraz-Schenkung, Julius-Klaus Stiftung and W.V. Scott Foundation, all to MK. SLK was supported by The Branco Weiss Fellowship - Society in Science.

Competing interests

The authors declare no competing interests.

Ethical statement

Permits for the use of animals for scientific purposes were granted by the Dept. of Biodiversity, Conservation and Attractions ([SF002958](#); [SF010888](#); [SF10388](#); [SF002958](#); [SF010774](#); [08-000920-1](#); [08-000706-3](#)) and the Dept. of Agriculture and Food ([U 10/2015-2018](#)). The animal ethics committees of the University of Western Australia, Murdoch University and the University of Zurich provided approvals for the ethical treatment of animals in scientific research ([R2649/14](#); [RA/3/100/1449](#); [RA/3/100/1464](#)).

References

1. Boyd R, Richerson PJ. 1995 Why culture is common, but cultural evolution is rare. *Proc. Br. Acad.* **88**, 77–93. (doi:citeulike-article-id:1339814)
2. van Schaik CP, Ancrenaz M, Borgen G, Galdikas B, Knott CD, Singleton I, Suzuki A, Utami SS, Merrill M. 2003 Orangutan cultures and the evolution of material culture. *Science* (80-.). **299**, 102–5. (doi:10.1126/science.1078004)
3. Whiten A, Goodall J, McGrew WC, Nishida T, Reynolds V, Sugiyama Y, Tutin CE, Wrangham RW, Boesch C. 1999 Cultures in chimpanzees. *Nature* **399**, 682–5. (doi:10.1038/21415)
4. Hoppitt WJE, Laland KN. 2013 *Social learning: an introduction to mechanisms, methods, and models*. Princeton and Oxford: Princeton University Press.
5. Krützen M, Mann J, Heithaus MR, Connor RC, Bejder L, Sherwin WB. 2005 Cultural transmission of tool use in bottlenose dolphins. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 8939–43. (doi:10.1073/pnas.0500232102)
6. Smolker R, Richards A, Connor R, Mann J, Berggren P. 1997 Sponge Carrying by Dolphins (Delphinidae, *Tursiops* sp.): A Foraging Specialization Involving Tool Use? *Ethology* **103**, 454–465. (doi:10.1111/j.1439-0310.1997.tb00160.x)
7. Kopps AM, Krützen M, Allen SJ, Bacher K, Sherwin WB. 2014 Characterizing the socially transmitted foraging tactic ‘sponging’ by bottlenose dolphins (*Tursiops* sp.) in the western gulf of Shark Bay, Western Australia. *Mar. Mammal Sci.* **30**, 847–863. (doi:10.1111/mms.12089)
8. Krützen M, van Schaik C, Whiten A. 2007 The animal cultures debate: response to Laland and Janik. *Trends Ecol. Evol.* **22**, 6; author reply 7. (doi:10.1016/j.tree.2006.10.011)
9. Laland KN, Janik VM. 2006 The animal cultures debate. *Trends Ecol. Evol.* **21**, 542–7. (doi:10.1016/j.tree.2006.06.005)
10. Laland K, Janik VM. 2007 Response to Krützen et al.: Further problems with the ‘method of exclusion’. *Trends Ecol. Evol.* **22**, 7. (doi:10.1016/j.tree.2006.10.011)
11. Krützen M, Willems EP, van Schaik CP. 2011 Culture and geographic variation in orangutan behavior. *Curr. Biol.* **21**, 1808–12. (doi:10.1016/j.cub.2011.09.017)
12. Franz M, Nunn CL. 2009 Network-based diffusion analysis: a new method for detecting social learning. *Proc. R. Soc. B* **276**, 1829–36. (doi:10.1098/rspb.2008.1824)

- 281 13. Hoppitt W, Boogert NJ, Laland KN. 2010 Detecting social transmission in networks. *J. Theor.*
282 *Biol.* **263**, 544–55. (doi:10.1016/j.jtbi.2010.01.004)
- 283 14. Allen J, Weinrich M, Hoppitt W, Rendell L. 2013 Network-based diffusion analysis reveals
284 cultural transmission of lobtail feeding in humpback whales. *Science (80-.).* **340**, 485–488.
285 (doi:10.1126/science.1231976)
- 286 15. Hobaiter C, Poisot T, Zuberbühler K, Hoppitt W, Gruber T. 2014 Social Network Analysis
287 Shows Direct Evidence for Social Transmission of Tool Use in Wild Chimpanzees. *PLoS Biol.* **12**,
288 e1001960. (doi:10.1371/journal.pbio.1001960)
- 289 16. Coussi-Korbel S, Fragaszy D. 1995 On the relation between social dynamics and social
290 learning. *Anim. Behav.* **50**, 1441–1453.
- 291 17. Farine DR, Aplin LM, Sheldon BC, Hoppitt W. 2015 Interspecific social networks promote
292 information transmission in wild songbirds. *Proc. R. Soc. B* **282**, 20142804.
293 (doi:10.1098/rspb.2014.2804)
- 294 18. Würsig B, Jefferson T. 1990 Methods of Photo-Identification for small Cetaceans. *Reports Int.*
295 *Whal. Comm.*
- 296 19. Krützen M, Barré L, Möller L, Heithaus M, Simms C, Sherwin W. 2002 A biopsy system for
297 small cetaceans: darting success and wound healing in *Tursiops* spp. *Mar. Mammal Sci.* **18**,
298 863–878.
- 299 20. Gilson A, Syvanen M, Levine K, Banks J. 1998 Deer gender determination by polymerase chain
300 reaction: validation study and application to tissues, bloodstains, and hair forensic samples
301 from California. *Calif. Fish Game* **84**, 159–169.
- 302 21. Wang J. 2011 COANCESTRY: a program for simulating, estimating and analysing relatedness
303 and inbreeding coefficients. *Mol. Ecol. Resour.* **11**, 141–5. (doi:10.1111/j.1755-
304 0998.2010.02885.x)
- 305 22. Wang J. 2007 Triadic IBD coefficients and applications to estimating pairwise relatedness.
306 *Genet. Res.* **89**, 135–53. (doi:10.1017/S0016672307008798)
- 307 23. Krützen M, Sherwin W, Berggren P, Gales N. 2004 Population structure in an inshore cetacean
308 revealed by microsatellite and mtDNA analysis: Bottlenose dolphins (*Tursiops* sp.) in Shark
309 Bay, Western Australia. *Mar. Mammal Sci.* **20**, 28–47.
- 310 24. Hoppitt WJE. 2018 NBDA: A Package For Implementing Network-Based Diffusion Analysis.

- 311 25. R Core Team. 2018 R: A language and environment for statistical computing. R Foundation for
312 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
- 313 26. Cairns SJ, Schwager SJ. 1987 A comparison of association indices. *Anim. Behav.* **35**, 1454–
314 1469.
- 315 27. Farine DR. 2013 Animal social network inference and permutations for ecologists in R using
316 asnipe. *Methods Ecol. Evol.* **4**, 1187–1194. (doi:10.1111/2041-210X.12121)
- 317 28. Calenge C. 2015 Home Range Estimation in R: the adehabitatHR Package. , 1–60.
318 (doi:10.1111/j.1365-2656.2006.01186.x)
- 319 29. Fieberg J, Kochanny CO. 2005 Quantifying home-range overlap: the importance of the
320 utilization distribution. *J. Wildl. Manage.* **69**, 1346–1359.
- 321 30. Wild S, Hoppitt W. 2018 Choosing a sensible cut-off point: assessing the impact of uncertainty
322 in a social network on the performance of NBDA. *Primates* (doi:10.1007/s10329-018-0693-4)
- 323 31. Mann J, Sargeant BL, Watson-Capps JJ, Gibson QA, Heithaus MR, Connor RC, Patterson E.
324 2008 Why do dolphins carry sponges? *PLoS One* **3**, e3868.
325 (doi:10.1371/journal.pone.0003868)
- 326 32. Sargeant BL, Wirsing AJ, Heithaus MR, Mann J. 2007 Can environmental heterogeneity
327 explain individual foraging variation in wild bottlenose dolphins (*Tursiops* sp.)? *Behav. Ecol.*
328 *Sociobiol.* **61**, 679–688. (doi:10.1007/s00265-006-0296-8)
- 329 33. Burnham K, Anderson D. 2002 *Model selection and multi-model inference: a practical*
330 *information-theoretic approach*. 2nd edn. New York: Springer.
- 331 34. Tyne J, Loneragan N, Kopps A, Allen S, Krützen M, Bejder L. 2012 Ecological characteristics
332 contribute to sponge distribution and tool use in bottlenose dolphins *Tursiops* sp. *Mar. Ecol.*
333 *Prog. Ser.* **444**, 143–153. See <http://www.int-res.com/abstracts/meps/v444/p143-153/>.
- 334 35. Mann J, Sargeant B. 2003 Like mother, like calf: the ontogeny of foraging traditions in wild
335 Indian ocean bottlenose dolphins (*Tursiops* sp.). In *The biology of traditions: models and*
336 *evidence* (eds D Fragaszy, S Perry), pp. 236–266. Cambridge, UK: Cambridge University Press.
- 337 36. Connor RC, Smolker RA, Richards AF. 1992 Two levels of alliance formation among male
338 bottlenose dolphins (*Tursiops* sp.). *Proc. Natl. Acad. Sci. U. S. A.* **89**, 987–90.
- 339 37. Connor RC, Krützen M. 2015 Male dolphin alliances in Shark Bay: Changing perspectives in a
340 30-year study. *Anim. Behav.* **103**, 223–235. (doi:10.1016/j.anbehav.2015.02.019)

38. Bizzozzero MR, Allen SJ, Gerber L, Wild S, King SL, Connor RC, Friedman WR, Wittwer S, Krützen M. 2019 Tool use and social homophily among male bottlenose dolphins. *Proc. R. Soc. B*, in press.
39. Patterson EM, Krzyszczyk E, Mann J. 2015 Age-specific foraging performance and reproduction in tool-using wild bottlenose dolphins. *Behav. Ecol.* **00**, 1–10. (doi:10.1093/beheco/arv164)
40. Gibson QA, Mann J. 2008 Early social development in wild bottlenose dolphins: sex differences, individual variation and maternal influence. *Anim. Behav.* **76**, 375–387. (doi:10.1016/j.anbehav.2008.01.021)
41. Krzyszczyk E, Patterson EM, Stanton MA, Mann J. 2017 The transition to independence: sex differences in social and behavioural development of wild bottlenose dolphins. *Anim. Behav.* **129**, 43–59. (doi:10.1016/j.anbehav.2017.04.011)
42. Zefferman MR. 2016 Mothers teach daughters because daughters teach granddaughters: The evolution of sex-biased transmission. *Behav. Ecol.* **27**, 1172–1181. (doi:10.1093/beheco/arw022)
43. Mann J, Patterson E. 2013 Tool use by aquatic animals. *Philos. Trans. R. Soc. London* **368**. (doi:http://dx.doi.org/10.1098/rstb.2012.0424)
44. Bacher K, Allen S, Lindholm AK, Bejder L, Krützen M. 2010 Genes or culture: are mitochondrial genes associated with tool use in bottlenose dolphins (*Tursiops* sp.)? *Behav. Genet.* **40**, 706–14. (doi:10.1007/s10519-010-9375-8)
45. Kopps AM, Ackermann CY, Sherwin WB, Allen SJ, Bejder L, Krützen M. 2014 Cultural transmission of tool use combined with habitat specializations leads to fine-scale genetic structure in bottlenose dolphins. *Proc. R. Soc. B* **281**. (doi:10.1098/rspb.2013.3245)
46. McElreath R, Strimling P. 2008 When Natural Selection Favors Imitation of Parents. *Curr. Anthropol.* **49**, 307–316. (doi:10.1086/524364)
47. Wild S, Allen SJ, Krützen M, King SL, Gerber L, Hoppitt WJE. 2019 Data from: Multi-Network-Based Diffusion Analysis reveals vertical cultural transmission of sponge tool use within dolphin matriline. *Dryad Digit. Repos.* (doi:doi:10.5061/dryad.sc26m6c)